

Nutrient dynamics associated with leaf litter decomposition of three agroforestry tree species (*Azadirachta indica*, *Dalbergia sissoo*, and *Melia azedarach*) of Bangladesh

Mahmood Hossain • Mohammad Raqibul Hasan Siddique • Md. Saidur Rahman
Md. Zaber Hossain • Md. Mahedi Hasan

Received: 2010-5-24 Accepted: 2010-10-27

© Northeast Forestry University and Springer-Verlag Berlin Heidelberg 2011

Abstract: *Azadirachta indica* A. Juss, *Dalbergia sissoo* Roxb., and *Melia azedarach* L. are little studied species in nutrient return capabilities from leaf litter decomposition to maintenance of the soil fertility despite their importance in agroforestry practices of Bangladesh. A leaf litter decomposition experiment was conducted using a litterbag technique to assess the nutrient return efficiency of these species. The decomposition rate of leaf litter was highest for *M. azedarach* and lowest for *D. sissoo*. Rainfall and temperature of study sites showed a significant ($p < 0.05$) positive relationship with the rate of leaf litter decomposition. The highest decay constant was observed for *M. azedarach* (6.67). Nitrogen and Phosphorus concentration in leaf litter showed a decreased trend sharply at the end of the first month, whereas rapid decrease of Potassium concentration was reported within 10 days. Conversely, higher concentration of nutrient was observed at the later stages of decomposition. All three species showed a similar pattern of nutrient release ($K > N > P$) during the decomposition process of leaf litter. Among the studied species, *D. sissoo* was best in terms of N and P return and *A. indica* was best in terms of K return.

Keywords: agroforestry; decay constant; decomposition; leaf litter and nutrient dynamics

Introduction

Depletion of nutrients and soil's organic matter is a serious threat to agricultural production and food security in many tropical countries (Lal 2004; Ajayi 2007). Such a decline leads to a decrease in agricultural productivity due to nutrient depletion and poor environmental quality (Lal 2004). Litter usually improves soil quality through the addition of organic matter, which enhances the soil's water holding capacity, water filtration, biodiversity, activity of soil microorganisms, and nutrient concentrations (Guo and Sims 1999; Bossa et al. 2005; Ngoran 2006). More than half of the nutrients taken up by plants return to the soil through several ways, among which decomposition of litter contributes the majority (Cole 1986) and the nutrient release patterns are related to climatic condition and litter quality (Khietwtam and Ramakrishnan 1993). Litter usually contains lower nutrients, compared to living plants (Toky and Ramakrishnana 1983; Sundarapandian and Swamy 1999). Among litter types, leaf litter contains comparatively higher concentrations of nutrients and returns the major source of nutrients to the soil (Meentemeyer et al. 1982; Anderson and Swift 1983; Liu et al. 2000). Amount of nutrients delivered by annual litterfall to the soil through decomposition is an important factor for sustainable forest production and provides an index of forest productivity (Swift et al. 1979; Attiwill and Leeper 1987; Guo and Sims 1999; Villela and Proctor 1999; Wardle 2002).

Agroforestry, a sustainable land-use system (Chundawat and Gautam 1993), and optimal production of any agroforestry practice is influenced by the combination of tree and agricultural crops, amount of soil nutrients, moisture conditions, and rate of organic matter decomposition (Nair 1984). It is believed that agroforestry promotes a more efficient cycling of nutrients than

Foundation project: This work is supported by Bangladesh Academy of Science and University Grants Commissions of Bangladesh.

The online version is available at <http://www.springerlink.com>

Mahmood Hossain (✉) • Mohammad Raqibul Hasan Siddique •
Md. Saidur Rahman

Forestry and Wood Technology Discipline, Khulna University, Khulna -
9208, Bangladesh. E-mail: mahmoodhossain@hotmail.com

Md. Zaber Hossain
Soil Science Discipline, Khulna University, Khulna-9208.

Md. Mahedi Hasan
Forestry and Wood Technology Discipline, Khulna University, Khulna-
9208.

Responsible editor: Zhu Hong

traditional agriculture system (Smiley and Kroschel 2010). *Azadirachta indica* A. Juss., *Dalbergia sissoo* Roxb., and *Melia azedarach* L. have been extensively planted in different agroforestry practices in Bangladesh (Baksha and Basak 2000) with combination of different crops (e.g. paddy, wheat, pineapple and vegetables) (Lodhiyal and Lodhiyal 2003). The selection of these tree species is usually guided by their morphological characteristics and local demand (Baksha and Basak 2000); however, the prime challenge of better management and sustainable production within agroforestry practices depends on the selection of tree species having efficient nutrient return capabilities (Khietwtam and Ramakrishnan 1993; Ibrahim et al. 2008) through the decomposition of litter which influences the nutrient cycling and formation of soil organic matter. Therefore, the aims of the present study were to examine and compare the decomposition and nutrient releasing pattern during the leaf litter degradation of *A. indica*, *D. sissoo*, and *M. azedarach*.

Materials and methods

Description of study site

This study was conducted in Khulna University campus in 2008 under mixed plantations of *A. indica*, *D. sissoo*, *Lagerstroemia speciosa*, *Emblia officinalis*, *Swietenia mahagoni*, *Leucaena leucocephala*, *Cassia siamea*, *M. azedarach* and so on. The tree density of the site is 800 stems ha⁻¹. The climate is humid-subtropical with a mean temperature of 18–23°C in winter and 27–31°C in summer. Mean annual rainfall is 1980 mm; summer (May to September) contributes about 81% of the annual rainfall while winter season contributes about 19% of rainfall. Soil is clayey and pH is around 7.9.

Sample collecting and processing

The leaf litter decomposition experiment was conducted using a litterbag technique (Mason 1977) for five months. Bunks of yellowish senescent leaves of *M. azedarach*, *A. indica* and *D. sissoo* were collected during the peak period of leaf shedding (March 2008). Leaf litter was air-dried at room temperature and mixed thoroughly. Two grams of leaf litter for individual species were taken as an individual sample. Individual samples were placed into a nylon bag of 300 mm×150 mm with 1-mm² mesh size. To prevent the leaves from folding and clumping they were laid flatly inside the bags. Eighty bags for each species were placed in the field and 10 bags were brought back to the laboratory for calculating fresh to oven-dry weight ratio at 80°C to constant weight.

Ten bags for each species were collected at ten days intervals for the initial month and subsequently at monthly intervals for the remaining months. The collected samples were then gently washed to remove sediments and dirt particles by using a soft brush under slowly running tap water followed by final rinsing in distilled water. Each sample was then oven-dried at 80 °C to constant weight.

Mass loss and decay constant

The loss in dry mass of samples was calculated from initial converted oven-dry weight and remaining mass. The rate of decomposition was calculated from percentage of mass loss divided by respective days of collection. Decay constants for leaf litter were calculated using negative exponential decay model.

$$X / X_0 = \exp^{-kt} \quad (\text{Olson 1963}) \quad (1)$$

where, X is the weight remaining at time t , X_0 the initial weight, \exp the base of natural logarithm, k the decay rate coefficient and t is the time in year.

Sample digestion and nutrients measurements in leaf litter

The oven-dried samples were processed according to Allen (1974). The processed leaf samples were weighted to 0.2 g and put into a digestion flask. Micro-kjeldahl digestion was carried out for the samples and the extract was filtered and diluted to 100 mL with distilled water (Allen 1974). Nitrogen (N) and Phosphorus (P) concentrations in sample extracts were measured according to Weatherburn (1967) and Parsons et al. (1984), respectively using UV-Visible Recording Spectrophotometer (SHIMADZU, UV-160A, Japan). Potassium concentration in sample extracts was measured by Atomic Absorption Spectrophotometer (PERKIN ELMER 4100, USA). The nutrient amounts released from leaf litters were calculated as differences between initial and final absolute amounts and also expressed as percentage of initial amounts.

Statistical analysis

The rates of mass loss and nutrients (N, P and K) concentrations in leaf litter were compared among the different stages of decomposition of the study and also with the tree species. This comparison was performed by two-way analysis of variance followed by Duncan Multiple Range Test (DMRT) using SAS 6.12 statistical software. Moreover, the relationships among the rate of leaf litter decomposition, monthly rainfall amount and monthly mean temperature were evaluated by correlation analysis using SAS 6.12 statistical software.

Results

Decomposition pattern

The mass was to the tune of 46%–63% in the three species at the end of the experiments (Fig. 1). Comparatively (ANOVA, $p < 0.05$), a rate of decomposition of *M. azedarach* leaf litter was higher, whereas leaf litter of *D. sissoo* showed a lower rate of decomposition (Fig. 2). Decay constants (k) were calculated for the whole experiment. The decay constant (k) was comparatively higher (6.67) for *M. azedarach*, followed by *A. indica* (5.12) and

D. sissoo (3.91). The rate of leaf litter decomposition of each species showed a significant ($p < 0.05$) positive relationship with monthly rainfall (range of r value is from 0.86 to 0.88.) and temperature (range of r value is from 0.71 to 0.81).




Fig. 1 Mass loss (%) of leaf litter of *Melia azedarach*, *Azadirachta indica* and *Dalbergia sissoo* during the decomposition process.




Fig. 2 Rate of decomposition (%) of leaf litter of *Melia azedarach*, *Azadirachta indica* and *Dalbergia sissoo* during the decomposition process.

Nutrients in leaf litter

The initial concentration of N and P in the leaf litter of *M. azedarach* was comparatively (ANOVA, $p < 0.05$) higher whereas a higher K concentration was observed in *A. indica* (Table 1). Rapid decrease in N and P concentration was observed at the end of first month, while rapid decrease in K concentration was reported within 10 days of the decomposition experiment. Conversely, after the rapid decrease, nutrient (N, P and K) concentration was increased at the later stages of decomposition of leaf litter. Moreover, leaf litter of all the species showed a similar pattern ($K > N > P$) in decreasing nutrient concentration throughout the decomposition process (Fig. 3–5). Comparatively higher amounts of N (about $0.03 \text{ kg} \cdot \text{t}^{-1}$) and P ($2.8 \text{ kg} \cdot \text{t}^{-1}$) were added to the soil from the leaf litter of *M. azedarach* whereas higher amount of K was added to the soil from *D. sissoo* leaf litter (Table 2).

Table 1. Initial concentration of nutrients in leaf litter of different species

Species	Nitrogen ($\mu\text{g} \cdot \text{g}^{-1}$)	Phosphorus ($\mu\text{g} \cdot \text{g}^{-1}$)	Potassium ($\mu\text{g} \cdot \text{g}^{-1}$)
<i>Melia azedarach</i>	35	3542	7268
<i>Azadirachta indica</i>	30	813	7455
<i>Dalbergia sissoo</i>	32	720	5464

Table 2. Amount of nutrients added from the leaf litter of different species

Species	Nitrogen ($\text{kg} \cdot \text{t}^{-1}$)	Phosphorus ($\text{kg} \cdot \text{t}^{-1}$)	Potassium ($\text{kg} \cdot \text{t}^{-1}$)
<i>Melia azedarach</i>	0.03	2.80	5.20
<i>Azadirachta indica</i>	0.02	0.59	6.04
<i>Dalbergia sissoo</i>	0.03	0.54	4.01




Fig. 3 Nitrogen concentration ($\mu\text{g} \cdot \text{g}^{-1}$) in leaf litter of *Melia azedarach*, *Azadirachta indica* and *Dalbergia sissoo* at different stages of decomposition




Fig. 4 Phosphorus concentration ($\mu\text{g} \cdot \text{g}^{-1}$) in leaf litter of *Melia azedarach*, *Azadirachta indica* and *Dalbergia sissoo* at different stages of decomposition

Discussion

Decomposition pattern

Comparatively a rate of decomposition of leaf litter was higher during the first 30 days, followed by a gradual mass loss for the subsequent 120 days (Fig. 2). This pattern of leaf litter decompo-

sition indicates two stages (Berg 1986; Takeda 1995; Semwal et al. 2003). The initial stage is typically met by a relatively larger decrease in mass which is usually attributed to leaching of readily soluble substances and non-lignified carbohydrates (Parsons et al. 1990; Ibrahima et al. 1995; Prescott 2005; Ibrahima et al. 2008). The further decrease in mass loss may be attributed to the release of a higher percentage of recalcitrant fractions like cellulose, lignin and tannin at the advanced stage of leaf litter decomposition (Bloomfield et al. 1993).




Fig. 5 Potassium concentration ($\mu\text{g}\cdot\text{g}^{-1}$) in leaf litter of *Melia azedarach*, *Azadirachta indica* and *Dalbergia sissoo* at different stages of decomposition

The studied species showed comparatively faster mass loss than other multipurpose tree species of the central Himalaya, India as reported by Semwal et al. (2003) and this variation in mass loss may be due to the litter quality and site factors (Heal et al. 1997). The decay constant varied from species to species and from temperate to tropical regions (Brinson 1990; Takeda 1996; Lockaby and Walbridge 1998; Torreta and Takeda 1999). Again, microclimatic variation in decay constant was also reported to be a factor within a given climatic region (Anderson and Swift 1983; Verhoef and Gunadi 2001). The decay constant range of this study ($k=6.67\text{--}3.91$) was found to be similar with the range of leaf litter decay of tropical trees as reported by Budelman (1988) and Palm and Sanchez (1990). The higher decay rate of leaf litter decomposition for *M. azedarach* could be an indicator of higher litter quality, compared to other studied species.

Nutrients in leaf litter

Comparatively higher initial concentration of N and P in leaf litter of *M. azedarach* and higher K concentration in leaf litter of *A. indica* indicated that capabilities of these species to retranslocate these nutrients were lower during the senescence of leaves (Berg and Laskowski 2006; Hagen-Thorn et al. 2006). Nutrients release from leaf litter follows a sequential lowering concentration in $\text{K} > \text{N} > \text{P}$, which is similar to the mobility pattern of these nutrients (Eaton et al. 1973; Waring and Schlesinger 1985). The faster decreasing of K concentration from leaf litter was observed as it is a non-structural element (Tisdale et al. 1993) and

also liable to be the most leachable cation during the decomposition of litter (Berg and Staaf 1987; Herra'ez and Ferna'ndez Marcos 2000; Guo and Sims 2002). A pattern of two distinct phases of decreasing nutrient concentration (N, P and K) was observed during the decomposition process (Fig. 3–5). At the first phase, an initial rapid decrease in nutrient (N, P and K) concentration was observed and the subsequent phase was immobilization of nutrients (Herra'ez and Ferna'ndez Marcos 2000; Guo and Sims 2002; Moore et al. 2006; Alvarez et al. 2008; Hough and Cole 2009). The initial rapid decrease of nutrients concentration may be ascribed to the loss of the soluble forms of nutrients at the initial stages of decomposition (Mahmood et al. 2007) and a slower release of nutrients at the later stages of litter decomposition governed by refractory components (Schlesinger 1985; Herra'ez and Ferna'ndez Marcos 2000). On the contrary, increased concentration of nutrients (N, P and K) at different stages of decomposition (Fig. 3–5) was attributed to microbial or non-microbial immobilization in the residual leaf litter, atmospheric deposition or the activity of fungi or heterotrophic organisms (Berg and Staaf 1987; Cameron and Spencer 1989).

Conclusion

The amount of nutrients added to the soil through leaf litter depends on the litter quality, climate, individual nutrient concentration and their dynamics during the process of decomposition. The added nutrients may contribute to the sustainability of soil fertility, which is becoming an important phenomenon for agroforestry practices. Among the considered species, *D. sissoo* was found to be the best in terms of N and P while *A. indica* was best in terms of K.

Acknowledgement

Authors wish to thank Forestry and Wood Technology Discipline, Khulna University, Khulna, Bangladesh for the logistic support throughout the study. Our special thanks also go to Mr. Mahbubul Karim, 180123 Albert Road, Leytonstone, London. UK E10 PA for the language editing of this manuscript.

References

- Ajayi OC. 2007. User acceptability of sustainable soil fertility technologies lessons from farmers knowledge, attitude, and practice in Southern Africa. *Journal of Sustainable Agriculture*, **30**: 21–40.
- Allen SE. 1974. *Chemical Analysis of Ecological Materials*. Oxford: Blackwell Scientific publication, pp. 1–165.
- Alvarez E, Marcos MLF, Torrado V, Sanjurjo MJF. 2008. Dynamics of macronutrients during the first stages of litter decomposition from forest species in a temperate area (Galicia, NW Spain). *Nutrient Cycling in Agroecosystem*, **80**: 243–256.
- Anderson JM, Swift MJ. 1983. Decomposition in tropical forest. In: S.L. Sutton, T.C., Whitmore and A.C. Chadwick (eds), *Tropical Rain forest: Ecology and Management*. Blackwell, Oxford: Special Publication Series of the British Ecological Society, pp 287–309, 498p.

- Attiwill PM, Leeper GW. 1987. *Forest Soils and Nutrient Cycles*. Melbourne: Melbourne University Press, pp. 202.
- Baksha MW, Basak AC. 2000. Mortality of sissoo (*Dalbergia sissoo* Roxb.) in Bangladesh. In: S., Appanah, G., Allard and S.M. Amaty (eds), *Die-Back of Sissoo (Dalbergia sissoo) Proceedings of International Seminar, Katmandu, Nepal*, 25–28 April 2000, FORSPA/FAO, Bangkok, pp. 1–4.
- Berg B, Laskowski R. 2006. *Litter decomposition: a guide to carbon and nutrient turnover*. Amsterdam: Elsevier Scientific Publishing, pp. 1–421.
- Berg B, Staaf H. 1987. Release of nutrients from decomposing white birch leaves and Scots pine needle litter. *Pedobiologia*, **30**: 55–63.
- Berg B. 1986. Nutrient release from litter and humus in coniferous forest soils—a mini review. *Scandinavian Journal of Forest Research*, **1**: 359–369.
- Bloomfield J, Vogt KA, Vogt DJ. 1993. Decay rate and substrate quality of fine roots and foliage of two tropical tree species in the Luquillo experimental forest, Puerto Rico. *Plant Soil*, **150**: 233–245.
- Bossa JR, Adams JF, Shannon DA, Mullins GL. 2005. Phosphorus and Potassium release from *Leucaena* leaves in three environment of Haiti. *Nutrient Cycling in Agroecosystem*, **73**: 25–35.
- Brinson M. 1990. Riverine forests. In: A.E., Lugo, M., Brinson and S. Brown (eds), *Forested wetlands. Ecosystems of the world 15*. The Netherlands, Amsterdam: Elsevier Scientific Publishing, pp 87–134.
- Budelman A. 1988. The decomposition of the leaf mulches of *Leucaena leucocephala*, *Glyricidia sepium* and *Flemingia macrophylla* under humid tropical conditions. *Agroforestry System*, **7**: 47–62.
- Cameron GN, Spencer SR. 1989. Rapid leaf decay and nutrient release in a Chinese tallow forest. *Oecologia*, **80**: 222–228.
- Chundawat BS, Gautam SK. 1993. *A Textbook of Agroforestry*. New Delhi: Oxford and IBH Publisher, pp. 64–69.
- Cole DW. 1986. Nutrient cycling in world forest. In: S.P. Gessel (ed), *Forest site and productivity*. The Netherlands: Martinus Nijhoff Publishers, pp. 104–115.
- Eaton JS, Likens GS, Bormann F. 1973. Throughfall and stemflow chemistry in a northern hardwood forest. *Journal of Ecology*, **61**: 495–508.
- Guo LB, Sims REH. 1999. Litter decomposition and nutrient release via litter decomposition in New Zealand eucalypt short rotation forests. *Agriculture Ecosystems and Environment*, **75**: 133–140.
- Guo LB, Sims REH. 2002. Eucalypt litter decomposition and nutrient release under a short rotation forest regime and effluent irrigation treatments in New Zealand. II. Internal effects. *Soil Biology and Biochemistry*, **34**: 913–922.
- Hagen-Thorn A, Varnagiryte I, Nihlgard B, Armolaitis K. 2006. Autumn nutrient resorption and losses in four deciduous forest tree species. *Forest Ecology and Management*, **228**: 33–39.
- Heal OW, Anderson JM, Swift MJ. 1997. Plant litter quality and decomposition: an historical overview. In: G. Cadisch and K.E. Giller (eds), *Driven by nature: plant litter quality and decomposition*. Willingford: CAB International, pp 3–30.
- Herra'ez Ortega L, Ferna'ndez Marcos ML. 2000. Materialization of eucalyptus debris in two sites in NW Spain. In: M. Madeira and P. Khanna (eds), *International symposium managing forest soils for sustainable productivity*. UTAD: Vila Real, Portugal, pp. 87–88.
- Hough Z, Cole CA. 2009. Aboveground decomposition dynamics in riparian depression and slope wetlands of central Pennsylvania. *Aquatic Ecology*, **43**: 335–349.
- Ibrahima A, Biyanzi P, Halima M. 2008. Changes in organic compounds during leaf litter leaching: laboratory experiment on eight plant species of the Sudano-guinea of Ngaoundere, Cameroon. *Forest*, **1**: 27–33.
- Ibrahima A, Joffre R, Gillon D. 1995. Changes in litter during the initial leaching phase—an experiment on the leaf-litter of Mediterranean species. *Soil Biology and Biochemistry*, **27**: 931–939.
- Khietwtam RS, Ramakrishnan PS. 1993. Litter and fine roots dynamics of a relict sacred grove forest at Cherrapunji in north-eastern India. *Forest Ecology and Management*, **60**: 327–344.
- Lal R. 2004. Soil carbon sequestration impacts on global climate change and food security. *Science*, **304**: 1623–1629.
- Liu WY, Fox JED, Xu ZF. 2000. Leaf litter decomposition of canopy trees, bamboo and moss in a montane moist evergreen broad-leaved forest on Ailao mountain, Yunnan, Southwest China. *Ecological Research*, **15**: 435–447.
- Lockaby BG, Walbridge MR. 1998. Biogeochemistry. In: M.G., Messina and W.H. Conner (eds), *Southern Forested Wetlands: Ecology and Management*. Boca Raton, FL: Lewis publishers, pp 149–172.
- Lodhiyal N, Lodhiyal LS. 2003. Biomass and net primary productivity of Bhabar Shisham Forests in Central Himalaya, India. *Forest Ecology and Management*, **176**: 217–235.
- Mahmood H, Saberi O, Misri K, Japar Sidik B. 2007. Nutrients dynamics associated with leaf litter degradation of *Bruguiera parviflora* (Whight and Arnold) at Kuala Selangor Mangrove forest, Malaysia. *Indian Journal of Forestry*, **30**: 325–330.
- Mason FC. 1977. *Decomposition*. London: The institute of biology's studies no. 74. Edward Arnold Limited, p. 58.
- Meentemeyer V, Box EO, Thompson R. 1982. World pattern and amounts of terrestrial plant litter production. *Bioscience*, **32**: 125–128.
- Moore TR, Trofymow JA, Prescott CE, Fyles J, Titus BD. 2006. Patterns of carbon, nitrogen and phosphorus dynamics in decomposing foliar litter in Canadian forests. *Ecosystems*, **9**: 46–62.
- Nair PKR. 1984. The role of trees in soil productivity and protection. In: P.K.R. Nair (ed), *Agroforestry Systems in the Tropics*. London: Kluwer Academic Publishers, pp. 567–589.
- Ngoran A, Zakra N, Ballo K, Kouame C, Zapta F, Hofman G, Cleemant OV. 2006. Litter decomposition of *Acacia auriculiformis* and *Acacia mangium* under coconut trees on quaternary sandy soils in Ivory Coast. *Biology and Fertility of Soils*, **43**: 102–106.
- Olson JS. 1963. Energy storage and the balance of producers and decomposers in ecological systems. *Ecology*, **44**: 322–331.
- Palm CA, Sanchez PA. 1990. Decomposition and nutrient release patterns of leaves of three tropical legumes. *Biotropica*, **22**: 330–338.
- Parsons TR, Yoshiki M, Carol ML. 1984. *A manual of Chemical and Biological Methods for Seawater Analysis*. New York: Pergamon Press, p. 173.
- Parsons WFJ, Taylor BR, Parkinson D. 1990. Decomposition of aspen (*Populus tremuloides*) leaf litter modified by leaching. *Canadian Journal of Forest Research*, **20**: 943–951.
- Prescott CE. 2005. Do rates of litter decomposition tell us anything we really need to know? *Forest Ecology and Management*, **220**: 66–74.
- Schlesinger WH. 1985. Decomposition of chaparral shrub foliage. *Ecology*, **66**: 1353–1359.
- Semwal RL, Maikhuri RK, Rao KS, Sen KK, Saxena KG. 2003. Leaf litter decomposition and nutrient release patterns of six multipurpose tree species of central Himalaya, India. *Biomass and Bioenergy*, **24**: 3–11.
- Smiley GL, Kroschel J. 2010. Yield development and nutrient dynamics in cocoa-gliricidiaagroforests of Central Sulawesi, Indonesia. *Agroforestry Systems*, **78**: 97–114.

- Sundarapandian SM, Swamy PS. 1999. Litter production and leaf litter decomposition of selected tree species in tropical forests at Kodayar in the Western Ghats, India. *Forest Ecology and Management*, **123**: 231–244.
- Swift MJ, Heal OW, Anderson JM. 1979. Decomposition in Terrestrial Ecosystems. Studies in Ecology 5. Oxford, UK: Blackwell Scientific Publications, p.372.
- Takeda H. 1995. A 5 year study of litter decomposition processes in a *Chamaecyparis obtusa*. *Forest Ecological Research*, **10**: 95–104.
- Takeda H. 1996. Templates for the organization of soil animals communities in tropical forests. In: I.M., Turner, C.H., Diong, S.S.L. Lim, P.K.L. Ng (eds), *Biodiversity and the Dynamics of Ecosystems*. DIWPA SERIES, Singapore, pp. 217–226.
- Tisdale SL, Nelson WL, Beaton JD, Havlin JL. 1993. Soil Fertility and Fertilizers. 5th eds. New Delhi: Prentice Hall of India Private Limited, pp 634.
- Toky OP, Ramakrishnan PS. 1983. Secondary succession following slash and burn agriculture in North-Eastern India. I. Biomass, litterfall and productivity. *Journal of Tropical Ecology*, **71**: 747–757.
- Torreta NK, Takeda H. 1999. Carbon and nitrogen dynamics of decomposing leaf litter in a tropical hill evergreen forest. *European Journal of Soil Biology*, **35**: 57–63.
- Verhoef HA, Gunadi B. 2001. Decomposition dynamics and nutrient flow in pine forest plantation in Central Java. In: M.V. Reddy (ed.), *Management of Tropical Plantation-Forests and Their Soil Litter System*. Science Publisher, Inc., Enfield, NH, USA, pp. 173–211.
- Villela DM, Proctor J. 1999. Litterfall mass, chemistry, and nutrient retranslocation in a monodominant forest on Maraca Island, Roraima, Brazil. *Biotropica*, **31**: 198–211.
- Wardle DA. 2002. Communities and Ecosystems: Linking the Aboveground and Belowground Components. Princeton, NJ: Princeton University Press, pp 138–140.
- Waring RH, Schlesinger WH. 1985. Forest ecosystems concepts and management. In: *Decomposition and Forest Soil Development*. New York: Academic Press Inc, p. 338.
- Weatherburn MW. 1967. Phenol-hypochlorite reaction for determination of ammonia. *Analytical Chemistry*, **39**: 971–974.